




A product of the National
Cooperative Soil Survey,
a joint effort of the United
States Department of
Agriculture and other
Federal agencies, State
agencies including the
Agricultural Experiment
Stations, and local
participants

Custom Soil Resource Report for Kittitas County Area, Washington

Beaver Creek Short Plat



January 12, 2010

Preface

Soil surveys contain information that affects land use planning in survey areas. They highlight soil limitations that affect various land uses and provide information about the properties of the soils in the survey areas. Soil surveys are designed for many different users, including farmers, ranchers, foresters, agronomists, urban planners, community officials, engineers, developers, builders, and home buyers. Also, conservationists, teachers, students, and specialists in recreation, waste disposal, and pollution control can use the surveys to help them understand, protect, or enhance the environment.

Various land use regulations of Federal, State, and local governments may impose special restrictions on land use or land treatment. Soil surveys identify soil properties that are used in making various land use or land treatment decisions. The information is intended to help the land users identify and reduce the effects of soil limitations on various land uses. The landowner or user is responsible for identifying and complying with existing laws and regulations.

Although soil survey information can be used for general farm, local, and wider area planning, onsite investigation is needed to supplement this information in some cases. Examples include soil quality assessments (<http://soils.usda.gov/sqi/>) and certain conservation and engineering applications. For more detailed information, contact your local USDA Service Center (<http://offices.sc.egov.usda.gov/locator/app?agency=nrcs>) or your NRCS State Soil Scientist (http://soils.usda.gov/contact/state_offices/).

Great differences in soil properties can occur within short distances. Some soils are seasonally wet or subject to flooding. Some are too unstable to be used as a foundation for buildings or roads. Clayey or wet soils are poorly suited to use as septic tank absorption fields. A high water table makes a soil poorly suited to basements or underground installations.

The National Cooperative Soil Survey is a joint effort of the United States Department of Agriculture and other Federal agencies, State agencies including the Agricultural Experiment Stations, and local agencies. The Natural Resources Conservation Service (NRCS) has leadership for the Federal part of the National Cooperative Soil Survey.

Information about soils is updated periodically. Updated information is available through the NRCS Soil Data Mart Web site or the NRCS Web Soil Survey. The Soil Data Mart is the data storage site for the official soil survey information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or a part of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means


for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD). To file a complaint of discrimination, write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue, S.W., Washington, D.C. 20250-9410 or call (800) 795-3272 (voice) or (202) 720-6382 (TDD). USDA is an equal opportunity provider and employer.

Contents

Preface	2
Soil Map	5
Soil Map.....	6
Legend.....	7
Map Unit Legend.....	8
Map Unit Descriptions.....	8
Kittitas County Area, Washington.....	10
201—Roslyn ashy sandy loam, 0 to 5 percent slopes.....	10
205—Xerofluvents, 0 to 5 percent slopes.....	11
237—Kladnick ashy sandy loam, 0 to 3 percent slopes.....	12
263—Volperie very paragravelly ashy sandy loam, 5 to 30 percent slopes.....	13

Soil Map

The soil map section includes the soil map for the defined area of interest, a list of soil map units on the map and extent of each map unit, and cartographic symbols displayed on the map. Also presented are various metadata about data used to produce the map, and a description of each soil map unit.



KLUTH
20-14-3300-0013

FORTUNE CREEK LLC
20-14-3300-0007

NIETSLH
20-14-3300-0008

JADOLE BIRGE LLC
20-14-3300-0009

COOL WATER LLC

SHEFFIELD
20-14-3300-0021

SHEFFIELD
20-14-3300-0022

SHEFFIELD
20-14-3300-0044

BACK COUNTRY RESOURCES
20-14-3300-0020

TEANAWAY RIDGE LLC
20-14-3200-0003

BACK COUNTRY RESOURCES
20-14-3300-0016

BACK COUNTRY RESOURCES
20-14-3300-0018

BACK COUNTRY RESOURCES
20-14-3300-0019

Sec 33

Sec 33

Sec 33

Sec 33


Sec 27

Sec 34


Sec 34



Custom Soil Resource Report Soil Map



Map Scale: 1:3,450 if printed on A size (8.5" x 11") sheet.



121° 6' 4"

MAP LEGEND

- Area of Interest (AOI)
- Area of Interest (AOI)
- Soils
- Soil Map Units
- Special Point Features**
 - Blowout
 - Borrow Pit
 - Clay Spot
 - Closed Depression
 - Gravel Pit
 - Gravelly Spot
 - Landfill
 - Lava Flow
 - Marsh or swamp
 - Mine or Quarry
 - Miscellaneous Water
 - Perennial Water
 - Rock Outcrop
 - Saline Spot
 - Sandy Spot
 - Severely Eroded Spot
 - Sinkhole
 - Slide or Slip
 - Sodic Spot
 - Spoil Area
 - Stony Spot

MAP INFORMATION

Map Scale: 1:3,440 if printed on A size (8.5" x 11") sheet.

The soil surveys that comprise your AOI were mapped at 1:24,000. Please rely on the bar scale on each map sheet for accurate map measurements.

Source of Map: Natural Resources Conservation Service
 Web Soil Survey URL: <http://websoilsurvey.nrcs.usda.gov>
 Coordinate System: UTM Zone 10N NAD83

This product is generated from the USDA-NRCS certified data as of the version date(s) listed below.

Soil Survey Area: Kittitas County Area, Washington
 Survey Area Data: Version 3, Jun 15, 2009

Date(s) aerial images were photographed: 7/27/2006

The orthophoto or other base map on which the soil lines were compiled and digitized probably differs from the background imagery displayed on these maps. As a result, some minor shifting of map unit boundaries may be evident.

Map Unit Legend

Kittitas County Area, Washington (WA637)			
Map Unit Symbol	Map Unit Name	Acres in AOI	Percent of AOI
201	Roslyn ashy sandy loam, 0 to 5 percent slopes	1.0	4.2%
205	Xerofluvents, 0 to 5 percent slopes	12.7	51.9%
237	Kladnick ashy sandy loam, 0 to 3 percent slopes	6.7	27.4%
263	Volperie very paragravelly ashy sandy loam, 5 to 30 percent slopes	4.0	16.4%
Totals for Area of Interest		24.5	100.0%

Map Unit Descriptions

The map units delineated on the detailed soil maps in a soil survey represent the soils or miscellaneous areas in the survey area. The map unit descriptions, along with the maps, can be used to determine the composition and properties of a unit.

A map unit delineation on a soil map represents an area dominated by one or more major kinds of soil or miscellaneous areas. A map unit is identified and named according to the taxonomic classification of the dominant soils. Within a taxonomic class there are precisely defined limits for the properties of the soils. On the landscape, however, the soils are natural phenomena, and they have the characteristic variability of all natural phenomena. Thus, the range of some observed properties may extend beyond the limits defined for a taxonomic class. Areas of soils of a single taxonomic class rarely, if ever, can be mapped without including areas of other taxonomic classes. Consequently, every map unit is made up of the soils or miscellaneous areas for which it is named and some minor components that belong to taxonomic classes other than those of the major soils.

Most minor soils have properties similar to those of the dominant soil or soils in the map unit, and thus they do not affect use and management. These are called noncontrasting, or similar, components. They may or may not be mentioned in a particular map unit description. Other minor components, however, have properties and behavioral characteristics divergent enough to affect use or to require different management. These are called contrasting, or dissimilar, components. They generally are in small areas and could not be mapped separately because of the scale used. Some small areas of strongly contrasting soils or miscellaneous areas are identified by a special symbol on the maps. If included in the database for a given area, the contrasting minor components are identified in the map unit descriptions along with some characteristics of each. A few areas of minor components may not have been observed, and consequently they are not mentioned in the descriptions, especially where the pattern was so complex that it was impractical to make enough observations to identify all the soils and miscellaneous areas on the landscape.

The presence of minor components in a map unit in no way diminishes the usefulness or accuracy of the data. The objective of mapping is not to delineate pure taxonomic classes but rather to separate the landscape into landforms or landform segments that have similar use and management requirements. The delineation of such segments

Custom Soil Resource Report

on the map provides sufficient information for the development of resource plans. If intensive use of small areas is planned, however, onsite investigation is needed to define and locate the soils and miscellaneous areas.

An identifying symbol precedes the map unit name in the map unit descriptions. Each description includes general facts about the unit and gives important soil properties and qualities.

Soils that have profiles that are almost alike make up a *soil series*. Except for differences in texture of the surface layer, all the soils of a series have major horizons that are similar in composition, thickness, and arrangement.

Soils of one series can differ in texture of the surface layer, slope, stoniness, salinity, degree of erosion, and other characteristics that affect their use. On the basis of such differences, a soil series is divided into *soil phases*. Most of the areas shown on the detailed soil maps are phases of soil series. The name of a soil phase commonly indicates a feature that affects use or management. For example, Alpha silt loam, 0 to 2 percent slopes, is a phase of the Alpha series.

Some map units are made up of two or more major soils or miscellaneous areas. These map units are complexes, associations, or undifferentiated groups.

A *complex* consists of two or more soils or miscellaneous areas in such an intricate pattern or in such small areas that they cannot be shown separately on the maps. The pattern and proportion of the soils or miscellaneous areas are somewhat similar in all areas. Alpha-Beta complex, 0 to 6 percent slopes, is an example.

An *association* is made up of two or more geographically associated soils or miscellaneous areas that are shown as one unit on the maps. Because of present or anticipated uses of the map units in the survey area, it was not considered practical or necessary to map the soils or miscellaneous areas separately. The pattern and relative proportion of the soils or miscellaneous areas are somewhat similar. Alpha-Beta association, 0 to 2 percent slopes, is an example.

An *undifferentiated group* is made up of two or more soils or miscellaneous areas that could be mapped individually but are mapped as one unit because similar interpretations can be made for use and management. The pattern and proportion of the soils or miscellaneous areas in a mapped area are not uniform. An area can be made up of only one of the major soils or miscellaneous areas, or it can be made up of all of them. Alpha and Beta soils, 0 to 2 percent slopes, is an example.

Some surveys include *miscellaneous areas*. Such areas have little or no soil material and support little or no vegetation. Rock outcrop is an example.

Kittitas County Area, Washington

201—Roslyn ashy sandy loam, 0 to 5 percent slopes

Map Unit Setting

Elevation: 1,900 to 2,400 feet
Mean annual precipitation: 30 to 40 inches
Mean annual air temperature: 43 to 45 degrees F
Frost-free period: 85 to 115 days

Map Unit Composition

Roslyn and similar soils: 85 percent
Minor components: 15 percent

Description of Roslyn

Setting

Landform: Terraces
Down-slope shape: Concave
Across-slope shape: Concave
Parent material: Glacial drift with a mantle of loess and volcanic ash

Properties and qualities

Slope: 0 to 5 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Well drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.57 to 1.98 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Available water capacity: Moderate (about 8.0 inches)

Interpretive groups

Land capability (nonirrigated): 3c
Other vegetative classification: grand fir/common snowberry/pinegrass (CWS336)

Typical profile

0 to 1 inches: Moderately decomposed plant material
1 to 8 inches: Ashy sandy loam
8 to 15 inches: Ashy sandy loam
15 to 37 inches: Loam
37 to 49 inches: Gravelly loam
49 to 60 inches: Gravelly loam

Minor Components

Nard

Percent of map unit: 10 percent

Volperie

Percent of map unit: 5 percent

205—Xerofluvents, 0 to 5 percent slopes

Map Unit Setting

Elevation: 500 to 2,500 feet

Mean annual precipitation: 7 to 50 inches

Mean annual air temperature: 43 to 50 degrees F

Frost-free period: 110 to 180 days

Map Unit Composition

Xerofluvents and similar soils: 85 percent

Minor components: 15 percent

Description of Xerofluvents

Setting

Landform: Flood plains, stream terraces

Down-slope shape: Concave

Across-slope shape: Concave

Parent material: Alluvium

Properties and qualities

Slope: 0 to 5 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Somewhat excessively drained

Capacity of the most limiting layer to transmit water (Ksat): High (1.98 to 5.95 in/hr)

Depth to water table: About 36 inches

Frequency of flooding: Frequent

Frequency of ponding: None

Available water capacity: Low (about 4.4 inches)

Interpretive groups

Land capability (nonirrigated): 4s

Other vegetative classification: Douglas-fir/elk sedge (CDG132)

Typical profile

0 to 2 inches: Moderately decomposed plant material

2 to 20 inches: Sandy loam

20 to 23 inches: Loamy sand

23 to 60 inches: Extremely cobbly sand

Minor Components

Racker

Percent of map unit: 10 percent

Aquolls

Percent of map unit: 5 percent

Landform: Flood plains

Ecological site: WET ALKALI MEADOW 6-9 PZ (R007XY603WA)

237—Kladnick ashy sandy loam, 0 to 3 percent slopes

Map Unit Setting

Elevation: 2,000 to 3,000 feet
Mean annual precipitation: 45 to 75 inches
Mean annual air temperature: 43 to 45 degrees F
Frost-free period: 90 to 120 days

Map Unit Composition

Kladnick and similar soils: 85 percent
Minor components: 15 percent

Description of Kladnick

Setting

Landform: Terraces
Down-slope shape: Concave
Across-slope shape: Concave
Parent material: Glacial outwash with a mantle of volcanic ash

Properties and qualities

Slope: 0 to 3 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Well drained
Capacity of the most limiting layer to transmit water (Ksat): High (1.98 to 5.95 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Available water capacity: Low (about 4.2 inches)

Interpretive groups

Land capability (nonirrigated): 6s
Other vegetative classification: grand fir/vine maple (CWS551)

Typical profile

0 to 1 inches: Moderately decomposed plant material
1 to 9 inches: Ashy sandy loam
9 to 15 inches: Gravelly ashy sandy loam
15 to 24 inches: Very gravelly sandy loam
24 to 60 inches: Extremely gravelly sand

Minor Components

Roslyn

Percent of map unit: 5 percent

Kachess

Percent of map unit: 5 percent

Bertolotti

Percent of map unit: 5 percent

263—Volperie very paragravelly ashy sandy loam, 5 to 30 percent slopes

Map Unit Setting

Elevation: 2,200 to 2,700 feet

Mean annual precipitation: 30 to 50 inches

Mean annual air temperature: 43 to 45 degrees F

Frost-free period: 80 to 110 days

Map Unit Composition

Volperie and similar soils: 80 percent

Minor components: 20 percent

Description of Volperie

Setting

Landform: Mountain slopes

Landform position (two-dimensional): Summit, shoulder, backslope

Down-slope shape: Linear

Across-slope shape: Convex

Parent material: Residuum from phyllite and schist with a mantle of volcanic ash

Properties and qualities

Slope: 5 to 30 percent

Depth to restrictive feature: 30 to 40 inches to paralithic bedrock

Drainage class: Well drained

Capacity of the most limiting layer to transmit water (Ksat): High (1.98 to 5.95 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None

Frequency of ponding: None

Available water capacity: Low (about 3.8 inches)

Interpretive groups

Land capability (nonirrigated): 4e

Other vegetative classification: grand fir/pinegrass (CWG124)

Typical profile

0 to 1 inches: Slightly decomposed plant material

1 to 8 inches: Very paragravelly ashy sandy loam

8 to 16 inches: Very paragravelly loam

16 to 38 inches: Gravelly loam

38 to 48 inches: Weathered bedrock

Minor Components

Nard

Percent of map unit: 5 percent

Roslyn

Percent of map unit: 5 percent

Custom Soil Resource Report

Kladnick

Percent of map unit: 5 percent

Bertolotti

Percent of map unit: 5 percent