

Ex 84

U.S. Geological Survey - Earthquake Hazards Program

Quaternary Fault and Fold Database of the United States

As of January 12, 2017, the USGS maintains a limited number of metadata fields that characterize the Quaternary faults and folds of the United States. For the most up-to-date information, please refer to the [interactive fault map](#).

Frenchman Hills structures, Frenchman Hills fault (Class A) No. 561a

Last Review Date: 2016-05-20

citation for this record: Lidke, D.J., compiler, 2003, Fault number 561a, Frenchman Hills structures, Frenchman Hills fault, in Quaternary fault and fold database of the United States: U.S. Geological Survey website, <https://earthquakes.usgs.gov/hazards/qfaults>, accessed 10/08/2021 07:21 PM.

Synopsis

General: The east-trending Frenchman Hills structures include the Frenchman Hills and Lind Coulee faults that show evidence suggestive of Quaternary offset (West and Shaffer, 1988 #5549; Shaffer and West, 1989 #5551; Geomatrix Consultants Inc., 1990 #5550). The Frenchman Hills anticline and related folds and some faults of the Frenchman Hills uplift, however, are only known to deform rocks of the Miocene Columbia River Basalt Group. Quaternary age growth or tightening of other folds in the Yakima fold belt, and perhaps of the Frenchman Hills folds, has been suggested and inferred from several local and regional geologic relations in the Yakima fold belt (Campbell and Bentley, 1981 #3513; Reidel, 1984 #5545; Reidel and others, 1994 #3539). Contemporaneous contraction across the region suggests that the Yakima folds are favorably oriented in the current strain field and accommodate the strain through active folding and possibly faulting (Pratt, 2012 #7397; Bjornstad and others, 2012 #7394 citing unpublished Zachariasen and others, 2006). As summarized by Bjornstad and others (2012 #7394), global positioning system (GPS) "data indicate relatively low (<1 mm/yr) but non-zero convergence

across the Yakima fold belt.... In general, these rates are higher than those calculated on Quaternary faults." Based on the growing consensus that the Frenchman Hills folds are cored by one or more buried Quaternary faults, the faults are reassigned to Class A as opposed to the prior Class B classification.

Sections: This fault has 3 sections. Sections defined here differ in lateral extent from the fault sources prescribed by Coppersmith and others (2014 #7402). Their western section is longer than section 561c.

Name comments

General:

Section: Refers to exposed and inferred faults and to fault-related features mapped along most of the north flank of the easterly trending Frenchman Hills and Frenchman Hills anticline (Grolier and Bingham, 1971 #5542; West and Shaffer, 1988 #5549; Geomatrix Consultants Inc., 1990 #5550; Schuster and others, 1997 #3760). These structural features are called the Frenchman Hills fault in Geomatrix Consultants Inc. (1990 #5550) and that name is used herein; earlier usage of this name is not known. The Frenchman Hills fault, as discussed by Geomatrix Inc. (1990 #5550) extends along the north flank of the Frenchman Hills from the Columbia River, eastward to just south of the southeast tip of Potholes Reservoir. The geologic map compiled by Schuster and others (1997 #3760), suggests that the fault may continue west of the Columbia River where it is exposed entirely in Miocene, Columbia River Basalt Group units. Quaternary offset along the fault west of the Columbia River has not been described or suggested and it is shown and included with the folds and other faults of the Frenchman Hills anticlinal uplift [561c].

County(s) and State(s)

GRANT COUNTY, WASHINGTON

Physiographic province(s)

COLUMBIA PLATEAU

Reliability of location

Good

Compiled at 1:100,000 and 1:250,000 scale.

Comments: Location of fault from GER_Seismogenic_WGS84 (<http://www.dnr.wa.gov/publications>)

/ger_portal_seismogenic_features.zip, downloaded 05/23/2016) attributed to 1:100,000-scale map by Gulick (1990 #5561), Reidel and Fecht (1994 #5565), and 1:250,000-scale map of Schuster and others (1997 #3760).

Geologic setting

The Frenchman Hills are in the northeastern part of the Yakima fold belt, a structural-tectonic sub province of the western Columbia Plateaus Province (Reidel and others, 1989 #5553; 1994 #3539). The Yakima fold belt consists of a series of generally east-trending narrow asymmetrical anticlinal ridges and broad synclinal valleys formed by folding of Miocene Columbia River basalt flows and sediments. In most parts of the belt the folds have a north vergence with the steep limb typically faulted by imbricate thrust faults. According to Reidel and others (1989 #5553) these frontal faults are typically associated with the areas of greatest structural relief. In the few places where erosion exposes the frontal faults deeper in the cores of the anticlinal ridges the faults are seen to become steeper with depth (as steep as 45–70°). Along their lengths the anticlines are commonly broken into segments ranging between 5 and 35 km long with boundaries defined by abrupt changes in fold geometry. Anticlinal ridges of the Yakima fold belt began to grow in Miocene time (about 16–17 Ma), concurrent with eruptions of Columbia River basalt flows, and continued during Pliocene time and may have continued to the present (Reidel and others, 1989 #5553; 1994 #3539).

The south-dipping Frenchman Hills and Lind Coulee faults are thrust faults that cut the north limbs of the Frenchman Hills anticline and Lind Coulee flexure, which are the principal folds of the Frenchman Hills anticlinal uplift. This uplift forms one of the many anticlinal ridges that comprise the Yakima fold belt in south-central Washington. The Frenchman Hills and Lind Coulee faults show evidence for Quaternary faulting events, but the folds and other faults of the Frenchman Hills uplift are only known to deform rocks of the Columbia River Basalt Group (Miocene).

Length (km)

This section is 51 km of a total fault length of 123 km.

Average strike

N87°W (for section) versus N83°W (for whole fault)

Sense of movement

Thrust

Comments: The Frenchman Hills fault is mapped as an exposed, but mostly buried, thrust fault (Grolier and Bingham, 1971 #5542; Gulick, 1990 #5561; Reidel and Fecht, 1994 #5565; Schuster and others, 1997 #3760).

Dip

20–60°

Comments: Geomatrix Consultants Inc. (1996 #4676) used fault dips of 30°, 45°, and 60° to estimate slip rates for the main fault underlying the Frenchman Hills. West and Shaffer (1988 #5549) and Shaffer and West (1989 #5551) reported dips of 21–40° for the Lind Coulee fault [561b], which is present directly east of the Frenchman Hills fault.

Paleoseismology studies**Geomorphic expression**

Just east of the Columbia River, the west end of the Frenchman Hills fault is exposed in Miocene basalt of the Columbia River Basalt Group (Grolier and Bingham, 1971 #5542; Geomatrix Consultants Inc., 1990 #5550; Reidel and Fecht, 1994 #5565; Schuster and others, 1997 #3760). Farther east, but still near the Columbia River, Grolier and Bingham (1971 #5542) show the fault juxtaposing Miocene rocks of the Columbia River Basalt Group on Quaternary sediments. More recent geologic maps by Reidel and Fecht (1994 #5565), Gulick (1990 #5561), and Schuster and others (Schuster and others, 1997 #3760) show these same sediments as Pliocene to Miocene sediments and do not show a fault at the locality where Grolier and Bingham (1971 #5542) showed the faulted Quaternary sediments. For this same locality Geomatrix Consultants Inc. (1990 #5550) reported that the locality had been greatly disturbed by agricultural activity after 1982, and they also reported that examination of 1982, color IR photographs of this locality showed numerous strongly expressed lineaments. The central and eastern parts of the Frenchman Hills fault are mapped as a buried thrust fault overlain by Quaternary loess deposits (Reidel and Fecht, 1994 #5565; Schuster and others, 1997 #3760). Geomatrix Consultants Inc. (1990 #5550) reported, and portrayed in their figure 3-5, numerous photogeologic lineaments along the central and eastern parts of the

fault; they indicated that these lineaments consisted of tonal contrasts and breaks in slopes.

Age of faulted surficial deposits

Geologic maps show Miocene rocks of the Columbia Plateau Basalt Group faulted along the west end of the fault and show the central and eastern parts of the fault buried by Pleistocene-Holocene loess deposits (Grolier and Bingham, 1971 #5542; Reidel and Fecht, 1994 #5565; Schuster and others, 1997 #3760). The geologic map by Grolier and Bingham (1978 #5543) also shows faulted Quaternary sediments along the western part of the fault, but more recent geologic maps (Gulick, 1990 #5561; Reidel and Fecht, 1994 #5565; Schuster and others, 1997 #3760) show these same sediments as unfaulted Miocene to Pliocene units. According to Geomatrix Consultants Inc. (1990 #5550), 1982 color IR photographs show numerous lineaments in these sediments along the western part of the fault, which are now obscured by post-1982 agricultural activity. Geomatrix Consultants Inc. (1990 #5550) reported, and portrayed in their figure 3-5, numerous photogeologic lineaments along the central and eastern parts of the fault, which are the parts of the fault that are shown as buried by Quaternary loess deposits on the geologic maps mentioned above. Geomatrix Consultants Inc. (1990 #5550) also reported that some of these lineaments are present in late Quaternary flood deposits and may suggest late Quaternary faulting, but they noted that they did not conduct detailed studies of these linear features.

Historic earthquake

Most recent prehistoric deformation

undifferentiated Quaternary (<1.6 Ma)

Comments: No definitive evidence of Quaternary displacement has been described along the Frenchman Hills fault. Studies by Geomatrix Consultants Inc. (1990 #5550) identified photogeologic lineaments in Pleistocene-Holocene loess deposits and late Quaternary flood deposits along what has been mapped as a buried trace of the fault. Geomatrix Consultants Inc. (1990 #5550) noted that the photogeologic lineaments they identified were not studied in detail, but they suggested some were probably fault-related and implied Quaternary, or perhaps late Quaternary, offset along the Frenchman Hills fault.

**Recurrence
interval**

Comments: Geomorphic or stratigraphic data needed for recurrence estimates directly related to the Frenchman Hills fault have not been reported. Piety and others (1990 #3733) used uplift rates calculated from 13.5 Ma volcanic rocks to estimate recurrence intervals of 1,220–61,100 years based on displacement per events of 0.02–1.0 m.

**Slip-rate
category**

Less than 0.2 mm/yr

Comments: No slip data in Quaternary deposits are available for the Frenchman Hills fault. Some data is available on uplift rates of Miocene volcanic rocks across the Frenchman Hills anticline and related folds. Piety and others (1990 #3733) report 221 m of uplift of 13.5 Ma volcanic rocks, which yield uplift rates of 0.02 mm/yr. Geomatrix Consultants Inc. (1995 #3593) used uplift of 200 m and horizontal offset of 300 m of 10.5 and 16.0 Ma volcanic rocks along estimated fault dips of 30°, 45°, and 60° to estimate long-term slip rates of 0.008–0.067 mm/yr for an inferred principle fault underlying the Frenchman Hills.

**Date and
Compiler(s)**

2003
David J. Lidke, U.S. Geological Survey

References

#7394 Bjornstad, B.N., Winsor, K., and Unwin, S.D., 2012, A summary of fault recurrence and strain rates in the vicinity of the Hanford site: Topical report prepared for the U.S. Department of Energy under contract DE-AC05-76RL01830, 90 p.

#3513 Campbell, N.P., and Bentley, R.D., 1981, Late Quaternary deformation of the Toppenish Ridge uplift in south-central Washington: *Geology*, v. 9, p. 519–524.

#5550 Geomatrix Consultants, Inc., 1990, Seismotectonic evaluation of the Walla Walla section of the Columbia Plateau geomorphic province for Grand Coulee, North, Dry Falls, Pinto, and O'Sullivan Dams; Soda Lake, north Scooteney, and south Scooteney dikes: Technical report to U.S. Department of the Interior, Bureau of Reclamation, Denver, Colorado, under Contract 6-CS-81-07310, April 1990, 129 p.

#3593 Geomatrix Consultants, Inc., 1995, Seismic design mapping, State of Oregon: Technical report to Oregon Department of Transportation, Salem, Oregon, under Contract 11688, January 1995, unpaginated, 5 pls., scale 1:1,250,000.

#4676 Geomatrix Consultants, Inc., 1996, Probabilistic seismic hazard analysis DOE Hanford site, Washington: Technical report to Westinghouse Hanford Company, Richland, Washington, under Contract WHC-SD-W236A-TI-002, Rev.1, February, 1996, 366 p.

#5542 Grolier, M.J., and Bingham, J.W., 1971, Geologic map and sections of parts of Grant, Adams and Franklin Counties, Washington: U.S. Geological Survey Geologic Investigations Map I-589, 1 sheet, scale 1:62,500.

#5543 Grolier, M.J., and Bingham, J.W., 1978, Geology of parts of Grant, Adams, and Franklin Counties, east-central Washington: Washington Division of Geology and Earth Resources Bulletin 71, 91 p.

#5561 Gulick, C.W., 1990, Geologic map of the Moses Lake 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 90-1, 9 p. pamphlet, 1 sheet, scale 1:100,000.

#3733 Piety, L.A., LaForge, R.C., and Foley, L.L., 1990, Seismic sources and maximum credible earthquakes for Cold Springs and McKay Dams, Umatilla Project, north-central Oregon: U.S. Bureau of Reclamation Seismotectonic Report 90-1, 62 p., 1 pl.

#7397 Pratt, T.L., 2012, Large-scale splay faults on a strike-slip fault system—The Yakima folds, Washington State: *Geochemistry, Geophysics, Geosystems*, v. 13, Q11004, doi: 10.1029/2012GC004405.

#5545 Reidel, S.P., 1984, The Saddle Mountains—The evolution of an anticline in the Yakima fold belt: *American Journal of Science*, v. 284, p. 942-978.

#5565 Reidel, S.P., and Fecht, K.R., 1994, Geologic map of the Priest Rapids 1:100,000 quadrangle, Washington: Washington Division of Geology and Earth Resources Open File Report 94-13, 22 p. pamphlet, 1

sheet, scale 1:100,000.

#3539 Reidel, S.P., Campbell, N.P., Fecht, K.R., and Lindsey, K.A., 1994, Late Cenozoic structure and stratigraphy of south-central Washington, *in* Lasmanis, R., and Cheney, E.S., eds., *Regional geology of Washington State*: Washington Division of Geology and Earth Resources, p. 159-180.

#5553 Reidel, S.P., Fecht, K.R., Hagood, M.C., and Tolan, T.L., 1989, The geologic evolution of the central Columbia Plateau, *in* Reidel, S.P., and Hooper, P.R., eds., *Volcanism and tectonism in the Columbia River flood-basalt province*: Geological Society of America Special Paper 239, p. 247-264.

#5554 Repasky, T.R., Campbell, N.P., and Busacca, A.J., 1998, Earthquake hazard study in the vicinity of Toppenish Basin, south-central Washington: Technical report to U.S. Geological Survey, Reston, Virginia, under Contract 1434-HQ-97-GR-03013, December 1998, 27 p., 7 pls.

#3760 Schuster, E.J., Gulick, C.W., Reidel, S.P., Fecht, K.R., and Zurenko, S., 1997, Geologic map of Washington-southeast quadrant: Washington Division of Geology and Earth Resources Geologic Map GM-45, 20 p. pamphlet, 2 sheets, scale 1:250,000.

#5551 Shaffer, M.E., and West, M.W., 1989, Quaternary faulting in the Frenchman Hills anticline, Yakima fold belt, central Columbia Basin, Washington: Geological Society of America Abstracts with Programs, v. 21, no. 5, p. 142.

#5549 West, M.W., and Shaffer, M.E., 1988, Interim draft report —Probabilistic and deterministic seismotectonic studies—O'Sullivan Dam and Potholes Reservoir, Washington: Technical report to U.S. Department of the Interior, Bureau of Reclamation, Denver, Colorado.